

BLOOD TRANSFUSIONS

Answers

To Your
Questions



Votre
gouvernement

Québec



Answers to Your Questions

This brochure is intended for people who may need a blood or blood product transfusion and for those who regularly receive transfusions. It provides answers to the most frequently asked questions.



Produced by:

**Direction des communications, ministère de la Santé
et des Services sociaux**

This document is available online and can be ordered at
www.msss.gouv.qc.ca by clicking **Publications**.

It may also be ordered at **diffusion@msss.gouv.qc.ca** or by mail at:

Ministère de la Santé et des Services sociaux
Direction des communications
Diffusion
Édifice Catherine-De Longpré
1075, chemin Sainte-Foy, 4^e étage
Québec (Québec) G1S 2M1

Masculine pronouns are used generically in this document.

Legal deposit

Bibliothèque et Archives nationales du Québec, 2019

Library and Archives Canada, 2019

ISBN 978-2-550-83248-5 (Print)

ISBN 978-2-550-83249-2 (PDF)

All rights reserved for all countries. Any reproduction whatsoever, translation or dissemination, in whole or in part, is prohibited unless authorized by Les Publications du Québec. However, reproduction or use for non-commercial personal, private study or scientific research purposes is permitted, provided the source is mentioned.

© Gouvernement du Québec, 2019

Are Blood Transfusions Safe?

Blood transfusions are very safe and, in some cases, may be the only way to save a life. The precautions taken to ensure the quality of blood products are increasingly reliable. Donors are selected based on very strict criteria and all blood donations undergo the most technologically advanced testing to detect known diseases and viruses.

All these measures have reduced the risk of disease transmission to very low levels. In Québec, the risk of being infected with AIDS or HIV (the human immunodeficiency virus) as a result of a transfusion is 1 in 23,100,000 transfused units. For hepatitis, which is a liver disease, the risk has been evaluated at 1 in 1,600,000 units for hepatitis B and 1 in 11,800,000 for hepatitis C¹. The risks are thus extremely low compared to the benefits of transfusions.

A compensation plan

If, despite these measures, a person receives a defective or contaminated blood product, the gouvernement du Québec will compensate that person if his or her health has been affected, regardless of who is responsible.

For more information on the Compensation plan for victims of a product distributed by Héma-Québec, please consult the website at **Québec.ca**, or call Services Québec at **1-855-881-9870 (toll free)**.

1. Héma-Québec. *Notice d'accompagnement portant sur les produits sanguins labiles*, décembre 2018, 56 pages (French only).

What Is a Blood Transfusion?

A blood transfusion is a treatment that involves giving blood or blood products to a person on the advice of a doctor.


What Is Blood?

Blood is the liquid circulating in the arteries and veins. It is essential for the body to function properly. Blood transports oxygen, nutrients that nourish the cells of the body, as well as other substances that help fight diseases.

Blood cells are formed in the bone marrow. Blood is made up of red blood cells, white blood cells, and platelets, which are suspended in a liquid called plasma. The adult human body contains an average of five to six litres of blood.

Depending on a person's state of health, one or more of these blood products may be required. Red blood cells, platelets, and plasma are the products most often transfused.

Blood and Blood Products



Plasma

Platelets

Red blood
cells

Plasma Derivatives

Clotting factors

Albumin

Immunoglobulins

Red Blood Cells

A drop of blood contains about five million red blood cells. The red colour is caused by an iron-rich protein in the red blood cells called hemoglobin, which carries oxygen. Red blood cells are transfused when a person has lost a lot of blood, for example as a result of an accident or major surgery. They may also be given to people suffering from anemia.

Red blood cells can be kept for 42 days and are stored at 1 to 6 °C.

White Blood Cells

White blood cells protect the body against bacteria and viruses. As soon as there is an infection, white cells begin to fight it. White blood cell transfusions are extremely rare.

White blood cells can be kept for no more than 24 hours and are stored at 20 to 24 °C.

Platelets

Platelets are smaller than red blood cells. Their most important role is to form blood clots to prevent or stop bleeding. Platelets are transfused when a person has suffered a severe loss of blood, when a person's platelets are not functioning properly, or when there are too few platelets circulating in the person's blood.

Platelets can be kept for up to 7 days and are stored at 20 to 24 °C.

Plasma

Plasma is yellowish in colour and is the liquid part of the blood. Plasma contains proteins that the body needs to function properly. Human blood consists of 55% plasma. Plasma transfusions are given to people with clotting factor deficiencies that could cause significant bleeding during surgery, for example.

Plasma can be kept frozen for up to one year.

Plasma Derivatives

Concentrated plasma derivatives are obtained using a process known as fractionation. They include clotting factors and proteins such as albumin and immunoglobulins. Plasma derivatives have many uses.

Generally, these products can be kept for over a year.




Where Do Blood and Blood Products Come From?

Most blood products mentioned in this brochure come from Québec donors through Héma-Québec or from donors in other parts of Canada through Canadian Blood Services. Only plasma used to produce derivatives may come from donors residing outside the country.

Not Everyone Can Be a Donor!

Blood donors are carefully selected. Only people who meet strict criteria can donate their blood.

People who want to give blood must first show a piece of I.D. and fill out a questionnaire about their state of health and other subjects related to the transmission of certain diseases.



This questionnaire is only one step in ensuring quality of our blood supply. Next, a small drop of blood is collected from the tip of a finger to check whether there is enough iron in the blood for the person to give blood.

Blood is always taken with new, sterile, disposable needles, bags, etc. that are used only once.



Donors Are Volunteers

In Québec and throughout Canada, people who give blood do so to help others. They are volunteers, which means they are not paid for their blood donations.

Giving blood means giving life. A single blood donation can help save four lives.

What Tests Are Done on Blood?

All blood donations are thoroughly tested for hepatitis B, hepatitis C, the AIDS virus or HIV (human immunodeficiency virus), syphilis, and the virus responsible for certain blood diseases (HTLV-I/II). Other tests, such as for West Nile virus, are also performed when necessary. **The tests are performed on the day the blood is collected. If the result of one of the tests is inconclusive or positive, the blood must be discarded.**

The blood is also analyzed to determine the blood type (A, B, AB, or O) and whether it is Rh positive (+) or Rh negative (-). At the hospital, before transfusions are given, several other tests are performed to ensure that the blood or blood product selected is the one best suited to the person who will receive the transfusion.

How Are Blood and Blood Products Transfused?

The method and the time it takes for a transfusion depend on the blood product being given and the health of the person receiving the transfusion. All the equipment used for the transfusion is new, sterile and disposable. It is only used once.

Red Blood Cells

Red blood cell transfusions take an average of 2 to 4 hours.

Platelets

Platelet transfusions take an average of 30 minutes to 2 hours.

Plasma

Plasma transfusions take an average of 1 to 2 hours.

Plasma Derivatives

The time needed to transfuse plasma derivatives depends on the type of product and the amount to be given.

What Are the Benefits of a Blood Transfusion?

Because of transfusions, many types of medical and surgical treatments can be provided. For example, transfusions are often given to treat premature babies, for heart surgery and organ transplants, for cancer and anemia treatments, and to people who have lost a lot of blood after an accident. In Québec, it is estimated that over 75,000 people receive blood or blood product transfusions each year.

What Are the Most Common Adverse Reactions?

Most transfusions pose no problem. Nurses carefully monitor people receiving blood transfusions to check for adverse reactions. However, despite all the precautions that are taken, a transfusion can sometimes cause an adverse reaction, some of which are described below.

Allergic Reactions

Transfusions cause allergic reactions in less than 1% of people receiving blood or blood products². These reactions can take the form of urticaria (hives) or other skin reactions that usually disappear with proper medication.

Fever

Transfused blood and blood products can also cause fever, with or without chills. Less than 1% of all transfusions cause this type of reaction, which can also be treated with appropriate medication³. In very rare cases, fever can be caused by bacterial contamination of the blood or blood product.



2. NAWAJ, K.I. et LAMBERT, G. *Les incidents et accidents transfusionnels signalés au système d'hémovigilance du Québec en 2012*, Institut national de santé publique, July 2015, 82 pages (French only).

3. *Ibid.*

Alloimmunization (Self-immunization)

After a transfusion, some people develop antibodies that can destroy their red blood cells. This complication, known as “alloimmunization”, generally does not cause any symptoms. However, it can be detected using a blood test. These people are very carefully monitored during future transfusions.

Other Reactions

Other reactions may occur. For example, a quick increase in the volume of liquid circulating in the veins during a transfusion may cause problems in the elderly or in people with heart problems.

In some cases, minor to severe breathing problems may occur.

Outpatients who receive transfusions should know that adverse reactions can occur once they return home. These adverse reactions can include skin irritation, fever, chills, jaundice (yellow skin), and back pain. If unusual symptoms do occur, they should immediately go to the hospital where the blood product was administered. If this is not possible, they can go to their local community services centre (CLSC), a medical clinic, or another health care facility.

Patients who have experienced an adverse reaction or abnormal symptom after a transfusion should always advise their doctor of this before receiving another transfusion.

Strict measures ensure that the right product is administered to the right person and that no errors occur that could cause serious complications. In addition, when taking a blood sample, and before each transfusion, medical staff must carefully check the name of the person receiving the transfusion to avoid errors.

Can a transfusion be replaced with another treatment?

Sometimes a blood or blood product transfusion can be avoided by using alternatives, such as the recovery of blood during surgery or the use of medication to correct anemia before surgery. The decision as to which treatment will be used will be made following a discussion with the doctor, based on the person's medical condition.

Informed Consent For a Blood Transfusion

As with any medical intervention or treatment, a doctor must obtain free and informed consent from patients, except during an emergency, before providing a transfusion.

Doctors are obliged to provide their patients with all relevant information about the proposed treatment and possible alternatives. They must be sure that patients understand the benefits and risks associated with transfusions and answer any questions they may have. The final decision is up to the patient.

A doctor can provide advice but cannot force a person to agree to a transfusion.





Blood Is a Source of Life

Transfusions Help Save Lives

**For more information
on transfusions,
visit the following websites:**

- Héma-Québec
www.hema-quebec.qc.ca
- Canadian Blood Services
www.blood.ca

Québec.ca

www.msss.gouv.qc.ca

**Santé
et Services sociaux**

Québec

