

# Quadrivalent Meningococcal Conjugate Vaccine

| June 2023 |

Vaccination,  
the best protection

By having your child vaccinated, you are protecting him or her against meningococcal infections and their complications. Meningitis (infection of the brain lining) and meningococemia (infection of the blood) are 2 serious infections caused by meningococcus. There are different serogroups (or types) of meningococcus, among other serogroups A, B, C, W and Y. This vaccine protects against serogroup A, C, W and Y.



Disease	Signs and symptoms	Possible complications
<b>Meningococcal infections</b>	<ul style="list-style-type: none"><li>• High fever</li><li>• Severe headache</li><li>• Nausea and vomiting</li><li>• General feeling of illness</li><li>• Red marks or tiny pin-size hemorrhages or bruises on the skin</li></ul>	<ul style="list-style-type: none"><li>• Deafness</li><li>• Scars</li><li>• Permanent brain damage</li><li>• Amputation of hands or feet</li><li>• Death (5 to 15 % of cases even after treatment)</li></ul>

# Information on the Quadrivalent Meningococcal Conjugate Vaccine

Vaccination is the protection against meningococcal serogroup A, C, W and Y infections and their complications. This vaccine is usually given in Secondary 3. The vaccine is recommended for persons with a high risk of infection from meningococcus. It is also recommended for persons travelling to regions where the risk of meningococcal infection is high.

**Some symptoms may be caused by the vaccine (e.g. redness at the injection site). Other problems may occur by chance and are not related to the vaccine (e.g. cold, gastro, headache).**

**Meningococcal quadrivalent conjugate vaccine is safe. Most reactions are harmless and do not last long.**

Frequency	Possible reactions to the vaccine	What to do
In most cases (more than 50% of people)	<ul style="list-style-type: none"> <li>• Pain at the injection site</li> </ul>	<ul style="list-style-type: none"> <li>• Apply a cold, damp compress at the injection site.</li> <li>• Use a medication for fever or discomfort if needed.</li> <li>• See a doctor if symptoms are severe.</li> </ul>
Very often (less than 50% of people)	<ul style="list-style-type: none"> <li>• Redness and swelling at the injection site</li> <li>• Headache, muscle pain, fatigue, discomfort, nausea, pain in joints, diarrhea, loss of appetite, irritability, drowsiness (sleepiness)</li> </ul>	
Often (less than 10 % of people)	<ul style="list-style-type: none"> <li>• Fever, shivering, vomiting, skin rash</li> </ul>	

We recommend that you stay at the clinic for at least 15 minutes after vaccination because allergic reactions may occur. If an allergic reaction occurs, the symptoms will appear a few minutes after the vaccination. The person giving the vaccine will be able to treat this reaction immediately.

**FOR ANY QUESTIONS,**  
contact the person giving the vaccine,  
or Info-Santé 811 or your doctor.

You can also consult  
**Québec.ca/vaccination**